

White-rot fungi in phenols, dyes and other xenobiotics treatment - a brief review

Tišma, Marina; Zelić, Bruno; Vasić-Rački, Đurđa

Source / Izvornik: **Croatian journal of food science and technology, 2010, 2, 34 - 47**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:109:059397>

Rights / Prava: [Attribution-NoDerivatives 4.0 International](#)/[Imenovanje-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-04-02**

REPOZITORIJ

PTF OS

PREHRAMBENO-TEHNOLOŠKI FAKULTET OSIJEK

dabar
DIGITALNI AKADEMSKI ARHIVI I REPOZITORIJ

Repository / Repozitorij:

[Repository of the Faculty of Food Technology Osijek](#)



White-rot fungi in phenols, dyes and other xenobiotics treatment – a brief review

Marina Tišma^{1*}, B. Zelić², Đurđa Vasić-Rački²

¹University of Josip Juraj Strossmayer in Osijek, Faculty of Food Technology Osijek, Franje Kuhača 20, 31000 Osijek, Croatia

²University of Zagreb, Faculty of Chemical Engineering and Technology, Marulićev trg 19, 10000 Zagreb, Croatia

review

Summary

Bioremediation is an attractive technology that utilizes the metabolic potential of microorganisms in order to clean up the environmental pollutants to the less hazardous or non-hazardous forms with less input of chemicals, energy and time. White-rot fungi are unique organisms that show the capacities of degrading and mineralizing lignin as well as organic, highly toxic and recalcitrant compounds. The key enzymes of their metabolism are extracellular lignolytic enzymes that enable fungi to tolerate a relatively high concentration of toxic substrates. This paper gives a brief review of many aspects concerning the application of white-rot fungi with the purpose of the industrial contaminants removal.

Keywords: bioremediation, lignolytic enzymes, white-rot fungi, waste treatment

Introduction

There is an increasing awareness around the world regarding the environmental pollution caused by industrial wastes. Billions of hazardous pollutants are produced annually by the chemical, agricultural, oil, paper, textile and other industries. Development of efficient, cost effective and sustainable methods as well as the improvement of the existing ones is becoming more and more important.

Bioremediation is an attractive technology that utilizes the metabolic potential of microorganisms in order to clean up the environmental pollutants to the less hazardous or non-hazardous forms with less input of chemicals, energy and time (Asgher et al., 2008, Haritash and Kaushik, 2009). Implementation of fungi in the process of remediation is called mycoremediation. Fungi play vital roles in all ecosystems, regulating the flow of nutrients and energy through their mycelial networks. It can be said that they act like natural and true ecosystem engineers (Singh, 2006). Great number of fungi, however, has not been discovered yet. There are data that approximately 80 000 to 120 000 species of fungi have been described to date, although the total number of these species is estimated at around 1.5 million (Webster and Weber, 2007). Fungi are known to degrade, or cause to deteriorate a wide variety of materials and compounds. They can degrade different type of wood, stored paper, textiles, plastics, leather and various wrapping materials. They can assist in deterioration of concrete or can cause decay of wall

paintings or can even attack ancient and medieval glass surfaces. Due to all of these facts, scientists have realized the possible benefits of the application of fungi in the complex areas in applied remediation engineering. Moreover, the discovery of the value of white-rot fungi in bioremediation has brought a great success in this field (Bennett, 2006; Singh, 2006). A significant progress has been achieved in the area of the white-rot fungi growth and enzyme production with the aim of enhancement the enzyme production. Molecular biology related to white-rot fungi, especially related to the extraction of genetic material (RNA and DNA), gene cloning and the construction of genetically engineered microorganisms is especially attractive and thus investigated in recent years (Gao et al., 2010.; Fan et al., 2010.). The present paper gives a brief review of the different application of white-rot fungi in the biodegradation of natural and artificial compounds presented in different industries with the aim to emphasize the importance of the biological waste treatment.

White-rot fungi

In comparison to bacteria most of fungi are robust organisms generally more tolerant to high concentrations of pollutants. It explains why they have been investigated extensively since the mid-1980s for their bioremediation capacities (Evans and Hedger, 2006; Strong, 2010; More et al., 2010). White-rot fungi (Fig. 1) constitute a diverse ecophysiological group

*Corresponding author: University of Josip Juraj Strossmayer in Osijek, Faculty of Food Technology Osijek, Franje Kuhača 20, 31000 Osijek, Croatia; Marina.Tisma@ptfos.hr

comprising mostly of basidiomycetes and litter-decomposing fungi (Wesenberg et al., 2003).



Fig. 1. White-rot fungi a) *Trametes versicolor* in the nature (Singh, 2006), b) *Trametes versicolor* G-99 in the laboratory, from spores to mycelial pellets

White-rot fungi possess a great range of different enzymes such as hydrolytic enzymes (cellulase, pectinase, xylanase) and extracellular ligninolytic enzymes (lignin peroxidases, manganese peroxidase and laccase) (Teerapatsakul et al., 2007). The expression pattern of these enzymes depends on the organism itself: some white-rot fungi produce lignin peroxidase and manganese peroxidase, but not laccase, while the others produce manganese peroxidase and laccase, but not lignin peroxidase (Hatakka, 1994). Therefore, among different types of white-rot fungi, some can equally decompose all of the three lignocellulose components in wood material, while some can degrade lignin and hemicellulose leaving cellulose intact (Lara et al., 2003; Bahri et al., 2006; Fang et al., 2008). Most of these enzymes are industrially important and have the great potential in processes of bioremediation,

biodegradation, biopulping, degradation and detoxification of recalcitrant substances (Wesenberg et al., 2003; Tortella et al., 2008).

In the nature, photosynthetic production of wood is balanced by degrading activity of different wood-destroying organisms (Leisola and Garcia, 1989). Wood is comprised of chemical substances called lignocellulose. Lignin, cellulose and hemicellulose are the main constituents of lignocellulose materials (Grönqvist et al., 2005; Mussatto et al., 2007). The complex architecture of wood is summarized in Fig. 2. Lignin is a three-dimensional biopolymer of high molecular weight. It is the most abundant aromatic compound on the earth and an important constituent of the living terrestrial biomass. Lignin is synthesized in plants by linking together hydroxycinnamyl, coniferyl and sinapyl alcohols to give *p*-hydroxyphenol (Bahri et al., 2006).




Fig. 2. Polymers in wood

White-rot fungi can degrade lignin in the way that the mycelia of the organisms penetrate the cell cavity and release lignolytic enzymes to decompose xylon to a white sponge-like mass (Gao et al., 2010). Beside lignolytic enzymes, in the process of lignin degradation many other enzymes are also involved: glyoxal oxidase, aryl oxidase, veratryl alcohol oxidase, oxalate decarboxylase, NAD-dependent formate dehydrogenase, P450 monooxygenase (Hattakka, 1994; Ander and Marzullo, 1997; Asgher, 2008). However, the main characteristic that differentiates white-rot fungi from most other microorganisms is their ability to completely mineralize all components of lignin to carbon dioxide and water (Asgher et al., 2008). Beside the role in wood degradation white-rot fungi can be applied for degradation of different industrial contaminants such as low molecular polycyclic aromatic carbohydrates, aromatic carbohydrates and chlorophenols (Harvey and Thurston, 2001; Cerniglica and Sutherland, 2001; Valentin et al., 2006; Valentin et al., 2007), textile dyes (Podgornik et al., 2001a; López et al., 2002; Wesenberg et al., 2003; Tychanowicz et al., 2004; Zille et al., 2005; Kariminiae-Hamedani et al., 2007) pesticides (Maloney, 2001) or in recent time pharmaceuticals such as ibuprofen, clofibrilic acid and carbamazepine (Marco-Urrea et al., 2009) or naproxen (Rodríguez-Rodríguez et al., 2010).

The ligninolytic systems have been widely studying in last few decades mainly on *Phanerochaete chrysosporium* (Leisola, 1984; Renganathan et al., 1990; Covert et al., 1992; Linko, 1992; Barclay et al., 1993; Jaspers and Penninckx, 1996; Rivela et al., 2000; Podgornik, 2001a; Podgornik, 2001b; Podgornik, 2002; Shahvali et al., 2000; Rodríguez Couto, 2002). In last two decades most of the investigations were done in order to find new varieties of white-rot fungi or to study the low cost production or overproduction of desired lignolytic enzymes. Furthermore, different investigations have been done in order to study the catalytic activity of produced enzymes on different substrates. According to the literature data, beside the *Phanerochaete chrysosporium*, some of the good producers of all lignolytic enzymes are also *Phanerochaete crassa* (Takano et al., 2006), *Pycnoporus cinnabarinus* (Sigoillot et al., 1999; Jonas et al., 2000; Geng and L., 2002; Alvarado et al., 2003; de Wilde, 2008; Gubta et al., 2010), *Pleurotus ostreatus* (Palmieri, 1997; Reddy et al., 2003; Membrillo et al., 2008) and *Bjerkandera adusta* (Moreira et al., 2001). Good laccase producers are *Trametes vesicolor* (Jang et al., 2002; Rancaño et al., 2003; Taveres et al., 2005; Revankar and Lele, 2006; Kurniawati and Nicell, 2007; Xavier et al.,

2007; Thiruchelvam and Ramsay, 2007; Tišma et al., 2008; Tišma et al., 2009; Tišma et al., 2010a; Tišma et al., 2010), *Pleurotis pulmonarius* (Tychanowicz et al., 2003), *Marasmius quercophilus* (Tagger et al., 1998) and *Polysporus ostreatus* (Claus, 2003).

The importance, properties and applications of lignolytic enzymes were described by several review papers (Dashtban et al., 2010; Lundell et al., 2010; Gianfreda et al., 1999; Mayer and Staples, 2002; Burton, 2003; Claus, 2003; Claus, 2004; Rodríguez Couto and Toca-Herrera, 2006; Rodríguez Couto and Toca-Herrera, 2007; Widsten and Kandelbauer, 2008; Linko, 1992; Klaus, 2002; Hamid and Rehman, 2009; Hofrichter, 2002; Hamid and Rehman, 2009).

A lot of research has been done in the field of genetic engineering, mainly on the most investigated fungus *Phanerochaete chrysosporium* (Kasai et al., 2010). There are several investigations where genes encoding ligninolytic enzymes in the white-rot fungi have been cloned and expressed in different hosts (Wang et al., 2004)

The potential of white-rot fungi can be harnessed thanks to emerging knowledge of the physiology and morphology (Žnidaršič and Pavko, 2001; Papagianni, 2004; Žnidaršič-Plazl, 2006) of these organisms. This knowledge could be transformed into reliable and robust waste treatment processes (Wesenberg et al., 2003).

In further text a brief overview of possible application of white-rot fungi and their enzymes in bioremediation purposes is given.

Application of white-rot fungi in treatment of phenolic compounds, dyes, and other xenobiotics

Numerous literature data are related to the treatment of different types of wastewater by fungi. The latest papers about degradation of phenolic compounds and specific industry wastewater treatment processes by white-rot fungi or their enzymatic system are presented in this review. Once released in the environment, pollutants can be biodegraded or bioaccumulated, non-transformed or block as soil-bound residue, and involved in non-biological transformation or physical processes (Gianfreda et al., 2006).

Removal of phenols and related compounds by white-rot fungi

Phenols are widely distributed compounds in the nature, especially in the plants (Boudet, 2007) where they occur in the form of alkaloids, coumarins, flavanoids, terpenes, tannins and lignins. They can

also be found in marine systems, produced by marine plants and animals where they can be degraded by indigenous microbial population. However, industrial production of phenols has been increasing. About 70 % of industrially produced phenols is employed in the production of resins and also used in the manufacture of plastic, biocides, disinfectants, textiles, medicines, explosives, inks, perfumes, photographic materials (Singh, 2006). Several types of industrial and agricultural wastes contain phenols (Gianfreda et al., 2003).

Pentachlorophenol (PCP) has been used since the late 1930s as a wood preservative together with its salt, sodium pentachlorophenate, due to its broad spectrum and low cost. Its esters have also been used as biocides. This generalized use has led to the contamination of many ecosystems. PCPs are currently considered as a priority product for decontamination studies according to the European Community and the American Environmental Protection Agency (Gomes Machado et al., 2005).

Huge amount of phenol-polluted waters are formed from the production of olive oil in Mediterranean region as well (Gianfreda et al., 2006). Olive mill wastewater (OMW), highly toxic effluent obtained from the extraction process by the olive oil industry, presents a major problem in the Mediterranean region. This effluent is variable in composition, but is always antibacterial and phytotoxic due to its phenolic content. The phenols responsible for the recalcitrant brownish color of OMWs are present in the residue as a mixture of monomeric aromatics and as polymerized heterogeneous pigments. The efficiency of dephenolization and decolorization of OMWs together with the production of laccase and manganese peroxidase were determined and compared by some authors showing that it can be effectively used as a pre-treatment step in a combined aerobic-anaerobic and/or physical and chemical treatment process to solve the environmental problems caused by OMW in Mediterranean olive oil-producing countries (Jaouani et al., 2003; Ergül et al., 2010).

Laconi et al. investigated the possibility of degradation of the high polyphenolic concentration of OMWs and, at the same time, they followed the production of microbial biomass that can be potentially useful as animal feed integrators. Four strains of the ligninolytic edible basidiomycete genus of *Pleurotus*, yeast strains *Saccharomyces cerevisiae* and *Kluyveromyces lactis*; the species of filamentous fungi *Oidodendron* spp. and *Penicillium* spp. were used in this study (Laconi et al., 2007).

Justino et al. reported the efficiency of three different approaches on phenols removal from OMW (Justino

et al., 2009; Justino et al., 2010). They oxidized different type of phenols that were extracted from OMW by a) biological treatment with two fungal species *Trametes versicolor* and *Pleurotus sajor*, b) enzymatic treatment by laccase, and c) chemical treatment by photo-Fenton oxidation. Phenols were removed more efficiently by photo-Fenton treatment than by biological or enzymatic treatments which indicates that additional investigations should be done in order to improve the biological approach.

D'Annibale et al. investigated the effect of submerged fermentation parameters, such as agitation and aeration, on growth and/or performance (i.e. lignin modifying enzymes production) of *Panus tigrinus*. The results of this study show that the up-scaling of OMWs treatment by *P. tigrinus* CBS 577.79 should be carried out in reactors providing good oxygen transfer with minimal shear effects such as the bubble column bioreactor. Moreover, due to the low aeration necessary for good mixing and mass-transfer the use of a simple bubble column bioreactor would have a positive impact on process costs (D'Annibale et al., 2006).

Phenolic compounds are part of coking wastewaters from the coke industry which are produced in the process of coking and coal liquefaction. Coking wastewaters contains complex inorganic and organic contaminants (ammonia, cyanide, thiocyanide, polycyclic nitrogen-containing aromatics, oxygen- and sulfur-containing heterocyclics and acyclic compounds) which are refractory, highly concentrated, toxic, mutative and carcinogenic (Lu et al., 2009). Lu et al. improved the process of phenolic compounds removal from the coking wastewater by the application of immobilized *Phanerochaete chrysosporium*.

Beside the possibility of the use of whole cells in the process of phenolic compound removal, considerable effort was done by different authors in the field of the application of pure enzymes. Bollag et al. successfully removed different chlorophenols presented alone or in combination of two or three by the pure laccase from *Trametes villosa* (Bollag et al., 2003.), D'Acunzo et al. removed different phenols by laccase with and without presence of different mediators (d'Acunzo et al., 2002). Lots of efforts are done on kinetics and modeling of phenol (Kurniawati and Nicell, 2006; Kurniawati and Nicell, 2007; Kurniawati and Nicell, 2008), catechol (Aktaş and Tanyolaç, 2003a; Aktaş and Tanyolaç, 2003b), pyrogallol (Güreşir et al., 2005), 1-naphtol (Aktaş et al., 2001) and L-DOPA (Tišma et al., 2008) oxidation catalyzed by laccase.

Improvements in the long-term application of enzymes and thereby a reduction in treatment costs,

could be also accomplished through the selection of an appropriate reactor configuration. Application of

white-rot fungi in the purpose of removal of phenols and related compounds is summarized in Table 1.

Table 1. White-rot fungi used in treatment of phenols and related compounds

White-rot fungi	Waste treated	References
<i>Pleurotus</i> sp.	Olive mill wastewater	Laconi et al., 2007
<i>Trametes versicolor</i> and <i>Pleurotus sajor</i>	Olive mill wastewater	Justino et al., 2009; Justino et al., 2010
<i>Panus tigrinus</i>	Olive mill wastewater	D'Annibale et al., 2006
<i>Phanerochaete chrysosporium</i>	Coking wastewaters	Lu et al., 2009
<i>Trametes villosa</i>	Wastewater containing chlorophenols	Bollag et al., 2003
<i>Poliporus pinsitus</i>	Oligomeric polyphenol compounds	D'Acunzo et al., 2002

Removal of polycyclic aromatic hydrocarbons (PAHs) by white-rot fungi

Polycyclic aromatic hydrocarbons (PAHs) are a class of ubiquitous xenobiotic environmental pollutants. The common sources of PAHs in environment include natural as well as anthropogenic. Natural sources are forest and rangeland fires, oil seeps, volcanic eruptions and exudates from trees. Anthropogenic sources of PAH include burning of fossil fuel, coal tar, wood, garbage, refuse, used lubricating oil and oil filters, municipal solid waste incineration and petroleum spills and discharge (Haritash and Kaushik, 2009). PAHs consist of three or more benzene rings fused in linear, angular or cluster arrangements which are thermodynamically stable. Based on the structure and mechanisms of activation, they exhibit toxic, mutagenic and carcinogenic properties. Improper disposal methods and inadequate control of these materials have created widespread contamination in soil groundwater and surface water. Bioremediation is shown to be effective for soils contaminated with low-molecular-weight PAHs (Singh, 2006). In last two decades the knowledge about the removal of PAHs by fungi is limited in comparison to that of bacteria. It is reported that some of white-rot fungi can efficiently degrade naphthalene, acenaphthene, anthracene, phenanthrene, fluorine, fluoranthene, chrysene, pyrene benzantracene, benzopyrene (Singh, 2006), chlorobenzene (Wang et al., 2008). Considerable efforts have been done with the aim of fungal bioremediation of different types of pesticides and polycyclic aromatic hydrocarbons by some authors (Eggen and Sveum, 1998; Eggen and Majcherczy, 1999; Valentín et al., 2006, Quintero et al., 2007; Acevedo et al., 2011). Clemente et al. investigated degradation of PAH by thirteen

deuteromycete ligninolytic fungal strains and found that the degree of degradation varies with a variation of lignolytic enzymes.

Maximum degradation of naphthalene (69 %) was observed by the strain 984 having manganese-peroxidase activity, followed by strain 870 (17 %) showing lignin peroxidase and laccase activities. Phenanthrene degradation of 12 % was observed with strain 870 with manganese-peroxidase and laccase activities. A good level of degradation of anthracene (65 %) was observed by the strain 710 (Clemente et al., 2001). Boyle et al. found that white-rot fungi growing in soil doesn't degrade significant amounts of PAHs. However, in liquid culture they degrade many PAHs (Boyle et al., 1998). The effect of nitrogen as nutrient was also assessed because nitrogen sources are frequently added during bioremediation. On the other hand nitrogen can inhibit the lignin-degrading system of white-rot fungi (Higson, 1991). Among hundreds of white-rot fungi displaying lignolytic activity, *Phanerochaete chrysosporium*, *Bjerkandera adusta* and *Pleurotus ostreatus* have been extensively studied. Intermediate compounds as quinones, hydroxyl- and dihydroxy-PAH have been isolated, but it is not clear whether they accumulate as dead-end products. Accumulation of PAH-quinones was reported in liquid cultures of *Phanerochaete chrysosporium* and *Bjerkandera adusta* and in soil by *Pleurotus ostreatus* (Haritash and Kaushik, 2009).

Application of white-rot fungi in the purpose of polycyclic aromatic hydrocarbons removal is summarized in Table 2.

Table 2. White-rot fungi used in treatment of polycyclic aromatic hydrocarbons

White-rot fungi	Waste treated	References
<i>Pleurotus ostreatus</i>	PAHs aged creosote contaminated soil	Eggen and Sveum, 1998; Eggen and Majcherczy, 1999
<i>Bjerkandera adusta</i> , <i>Irpex lacteus</i> and <i>Lentinus tigrinus</i>	PAHs in forest and salt marsh soils	Valentín et al., 2006
<i>Bjerkandera adusta</i>	hexachlorocyclohexane (HCH) isomers present in a spiked soil	Quintero et al., 2007
<i>Phanerochaete chrysosporium</i> and <i>Pleurotus pulmonarius</i>	aromatic hydrocarbons in an aged contaminated soil containing high concentrations of heavy metals	Boyle et al., 1998
<i>Phanerochaete chrysosporium</i>	xenobiotics in soil	Higson, 1991

Decolorization of dyes by white-rot fungi

The textile industry is the most frequent user of synthetic dyes. Dyes can be classified according to their structure (particularly the nature of the chromophore) or the method of application (Knapp et al., 2006). *Phanerochaete chrysosporium* was the first identified fungus able to degrade polymeric synthetic dyes. Up to date most investigations on dyes degradation have been performed by this fungus (Shahvali et al., 2000; Podgornik et al., 2001a; Singh et al., 2009; Singh and Pakshirajan, 2010; Nilsson et al., 2006; Faraco et al., 2009; Lucas et al., 2008; Levin et al., 2010). However it has been shown that it was not the best one (Lucas et al., 2008). A lot of efforts were done with some other species such as *Coriolus versicolor* (Hai et al., 2006; Nilsson et al., 2006; Erkurt et al., 2007; Asgher, 2009), *Pleurotus ostreatus* (Nilsson et al., 2006; Erkurt et al., 2007; Faraco et al., 2009; Lu et al., 2008) *Trametes trogii* and *Trametes villosa* (Levin et al., 2008; Levin et al., 2010). It has been proved that the mechanism of the oxidative enzymes in the dyes decolorization is strain dependant. For a fast screening of numerous fungal strains for their ability to decolourise textile dyes, a microtitre plate-based method was developed recently (Lucas et al., 2008).

Vijaykumar et al. isolated a novel fungus *Cladosporium cladosporioides* from coal sample and used it in the process of decolorization of five different azo and triphenylmethane dyes like acid blue 193, acid black 210, crystal violet, reactive black B(S) and reactive black BL/LPR both on solid and in liquid broth medium (Vijaykumar et al., 2006). Azo dyes such as DR-80 are important colorants and constitute the largest class of dyes for application not only in textile, but also in paper, leather, gasoline, foodstuffs and cosmetics industries (Sing et al., 2009). Shahvali et al. investigated the effect of different parameters (size of inoculum, temperature,

carbon source) on decolorization of textile wastewaters using *Phanerochaete chrysosporium*. *Phanerochaete chrysosporium* was able to decolorize textile effluents with efficiency of up to 97 % (Shahvali et al., 2000). With the aim of decolorization of cotton bleaching effluent, Zhang et al. successfully developed a continuous fluidized-bed bioreactor (Zhang et al., 1998).

Singh et al. investigated decolourization of Direct Red 80 (DR-80) with *Phanerochaete chrysosporium* MTCC 787 by employing sequential design of experiments. Media components for growing the white-rot fungus were first screened using Plackett-Burman design and then optimized using response surface methodology (RSM), which resulted in enhancement in the efficiency of dye removal by the fungus. At the RSM optimized levels of the media constituents, *Phanerochaete chrysosporium* showed complete (100 %) dye decolorization efficiency due to its maximum LiP activity (Sing et al., 2009).

Nilsson et al. investigated the removal of two dyes, Reactive Blue 4 and Reactive Red 2 by different white-rot fungi *Phanerochaete chrysosporium*, *Trametes versicolor*, *Pleurotus ostreatus* and *Pleurotus sajor-caju*, in a reactor systems with immobilized mycelia (Nilsson et al., 2006).

Capabilities of the fungi *Phanerochaete chrysosporium* and *Phanerochaete ostreatus* and of free and immobilized laccase mixtures from *Phanerochaete ostreatus* on industrial dye wastewaters have been demonstrated by Faraco et al. Wastewater model containing dyes with complex trisazo-, polyazo- and stilbene- type structures was decolorized by *Phanerochaete chrysosporium* (about 45 % decolourization in only 1 day of treatment). The acid wastewater model was decolorized by *Phanerochaete ostreatus* (60 % decolorization in 7 days). Based on the discharged amounts, economic relevance and representativeness of chemical structures of the contained dyes, models of acid,

direct and reactive dye wastewaters from textile industry have been defined (Faraco et al., 2009). Decolourization of two azo dyes, Direct Red-80 and Mordant Blue-9 by *Phanerochaete chrysosporium* was investigated both individually and in mixtures in batch shake flasks. The profile of enzyme activities and dye decolourization by the fungus, in both single and mixed dye systems, suggested that fungal peroxidase enzymes (MnP and LiP) play a strong role in these dyes decolourization (Singh and Pakshirajan, 2010). Chander and Arrora have shown that

Dichomitus squalens, *Daedalea flavida*, *Irpex flavus* and *Polyporus sanguineus* are better decolorizers of laboratory dyes than the much studied *Phanerochaete chrysosporium*. The fungal-based biocleaning systems have been suffering from drawback of adsorption, thus, in order to overcome this limitation, the cell free enzyme extracts obtained from fungal cultures have been used (Chander and Arrora, 2007). Application of white-rot fungi in the purpose of different dyes removal is summarized in Table 3.

Table 3. White-rot fungi used in treatment of dyes

White-rot fungi	Dye(s)	References
<i>Phanerochaete chrysosporium</i>	Anthraquinone dyes	Lucas et al., 2008
<i>Phanerochaete chrysosporium</i>	Azo dye	Shahvali et al., 2000
<i>Phanerochaete chrysosporium</i>	Indigo carmine	Podgornik et al., 2001a
<i>Phanerochaete chrysosporium</i>	Direct Red-80	Sing et al., 2009
<i>Phanerochaete chrysosporium</i> , <i>Coriolus versicolor</i> , <i>Pleurotus ostreatus</i> and <i>Pleurotus sajor-caju</i>	Reactive Blue 4 and Reactive Red 2	Nilsson et al., 2006
<i>Phanerochaete chrysosporium</i> and <i>Phanerochaete ostreatus</i>	Direct Blu 71, Direct Red 80, Direct Yellow 106, Reactive Blue 222, Reactive Red 195, Reactive Yellow 145, Reactive Black 5, Acid Blue 62, Acid Yellow 49, Acid Red 266	Faraco et al., 2009
<i>Coriolus versicolor</i>	A synthetic waste water with Poly S119 dye	Hai et al., 2006
<i>Pleurotus ostreatus</i> , <i>Coriolus versicolor</i> and <i>Funalia trogii</i>	Remazol Brilliant Blue Royal and Drimaren Blue	Erkurt et al., 2007
<i>Coriolus versicolor</i>	Crescent Textile Industry (CRT), Itmad Textile Industry (ITT), Megna Textile Industry (MGT) and Ayesha Textile Industry (AST) effluents	Asgher, 2009
<i>Pleurotus ostreatus</i>	Acid Orange 7, Acid Orange 8 and Mordant Violet 5	Lu et al., 2008
<i>Cladosporium cladosporioides</i>	Acid blue 193, acid black 210, crystal violet, reactive black B(S) and reactive black BL/LPR	Vijaykumar et al., 2006
<i>Dichomitus squalens</i> , <i>Daedalea flavida</i> , <i>Irpex flavus</i> and <i>Polyporus sanguineus</i>	Coracryl dyes (black, pink, violet, red) Reactive dyes (yellow, orange and red) Rathiodal dyes (scarlet)	Chander and Arrora, 2007

Decolorization of industrial effluents by white-rot fungi

The application of white-rot fungi in large-scale waste treatment, however, has been impeded owing to the lack of an appropriate reactor system capable of coping with rather slow fungal degradation, loss of the extracellular enzymes and mediators with discharged water, and excessive growth of fungi. In this context, a feasible system may be envisaged by coupling the excellent degradation capability of the white-rot fungi with the inherent advantages of a membrane bioreactor (MBR), yielding reduced excess sludge production.

Accomplishment of excellent stable pollutant removal (99 % color and 97 % TOC removal), using *Coriolus versicolor* along with the alleviation of the membrane fouling problem by employing a reasonable chemical cleaning dose is however a novel and attractive system (Hai et al., 2006).

Asgher et al. in their work presented the screening of *Coriolus versicolor* IBL-04 on five effluents of different industries. Optimization of different process parameters for Arzoo Textile Industry (ART) effluent decolorization showed that manganese peroxidase (MnP) was the lignolytic enzyme present in the culture filtrates, while lignin peroxidase (LiP) and laccase were undetectable

(Asgher et al., 2009). From the other hand, Erkurt et al. showed that laccase was the only enzyme responsible for decolorization of Remazol Brilliant Blue Royal (RBBR) and Drimaren Blue CL-BR by *Pleurotus ostreatus*, *Coriolus versicolor* and *Funalia trogii* (Erkurt et al., 2007).

Three sulphonated phenylazonaphthol dyes with similar molecular structures, Acid Orange 7, Acid Orange 8 and Mordant Violet 5 were selected and degraded by the white-rot fungus *Pleurotus ostreatus* (Lu et al., 2008).

The pulp and paper industry is one of the primary users of wood resources in the world. Pulp and paper industries produce annually over 2800 billions' lives of colored, toxic and intensely colored waste effluents, causing severe water pollution wastewater (Jaspers and Penninckx, 1996). This kind of effluent, usually called black liquor, has a high level of chemical oxygen demand. The primary contributors to the color and toxicity of these effluents are high-molecular-weight lignin and its derivatives (Wu et al., 2004). No information on the exact chemical nature of the chromophores in bleaching effluent exists. However it is likely that the chromophores are different to some extent as wood has a very high lignin content (which is removed during the treatment) and cotton does not have it. In cotton bleaching, absorbable organically bound halogens are produced when chlorine is used in the process (Zhang et al., 1998). Although currently available methods, such as chemical oxidation, reverse osmosis and adsorption, are highly efficient, they suffer some disadvantages. The limitations include high cost, limited applicability, high energy input, and usually these treatments may result in the production of toxic by-products (Hamid and Rehman, 2009).

Over the past 20 years, pollution caused by pulp black liquor has been seriously increased. Black liquor contributes only 10–15 % of the total wastewater, but accounts for nearly 80 % of color, 30 % of the biochemical oxygen demand and 60 % of the chemical oxygen demand of the total pollution load of pulp and paper mill effluent.

Different approaches have been used in order to solve this problem. Shararri et al. used bagasse effluent collected from the wastewater collection tank from pulp and paper industry for the biotreatment with *Phanerochaete chrysosporium* (Shararri et al., 2010). Jaspers and Penninckx investigated a possible adsorption of color and toxic chloroderivatives of lignin absorbable organically bound halogens that are essentially formed in bleaching process (the Kraft process) when using preformed pellets for inoculation. They showed that depending of the

conditions of incubation, the pellets of the fungus can strongly absorb color and absorbable organically bound halogens from the Kraft beach plant effluent (Jaspers and Penninckx, 1996). A repeated batch operation is developed for the treatment of alkaline pulp black liquor, through a process of biological acidification and precipitation of lignin using brown-rot fungus *Fomitopsis* sp. IMER2 (Ma et al., 2008).

Wu et al. used individually different white-rot fungi, *Phanerochaete chrysosporium*, *Pleurotus ostreatus*, *Lentinus edodes*, *Trametes versicolor* and strain S22 grown on a porous plastic media to treat black liquor from a pulp and paper mill. Three white-rot fungi, *Phanerochaete chrysosporium*, *Pleurotus ostreatus* and S22, showed high capacity for lignin degradation at pH 8.0-11.0, suggesting that the white-rot fungi were able to grow and degrade lignin even under strong alkaline conditions (Wu et al., 2005).

Distillery wastewater is produced as a result of distillation of ethanol produced in the fermentation of carbohydrates. For every liter of produced ethanol, up to 20 L of stillage is generated. Therefore, in a typically distillery, over a half million liters of stillage is generated every day. The composition of molasses stillage depends on the source of raw material used for the alcohol production. For an example, the composition of sugar beet and sugarcane molasses stillage includes various organic compounds, like acetic acid, lactic acid, glycerol and reducing sugars.

Alcohol distillery wastewater (ADW) is a dark brown colored wastewater whose color components disrupt the flow of penetration of sun rays in surface waters, which in turn reduces the photosynthetic activity and is detrimental to aquatic life. The ADW water is recalcitrant due to the presence of melanoides. ADW has antioxidant properties and decolorization of ADW is also recalcitrant to normal biological wastewater treatments. Certain physical, chemical and thermal methods have been used but they are unsuitable for commercial application. On the other hand highly suitable treatment, such as anaerobic digestion, involves high installation and implementation costs.

In the 1990s, several researchers used white-rot fungi in the decolorization of distillery effluent. It appeared that decolorization was attributed to the secretion of extracellular lignolytic enzymes. The advantage of the use of white-rot fungi, beside the decolorization, is production of by-product – a single-cell protein (Singh, 2006). In last decades, different approaches have been used. Pant and Adholeya investigated the production of lignolytic enzymes by fungi isolated from distillery effluent and effluent contaminated soils by cultivation of the isolated fungi on wheat straw and corncob powder as substrates. Among all

isolated species, maximum decolorization of effluents from a cane molasses based distillery (86.33 %) was achieved by *Pleurotus ostreatus* (Pant and Adholeya, 2007).

Many scientists have been working in the field of isolation of the new strains. Chairattananokorn et al. found new *Pycnoporus coccineus* strains FPF 00062506, FPF 97062901, FPF 97091303, and FPF 98063001 that have the potential for ADW decolorization at thermophilic conditions (Chairattananokorn et al., 2005).

Strong recently published a paper where wastewater from distillation of fermented marula fruit was treated by four white-rot fungi (*Trametes*

pubescens MB 89, *Ceriporiopsis subvermispora*, *Pycnoporus cinnabarinus* and *Phanerochaete chrysosporium*) in shake flasks experiments at pH 5.0 with no additional carbon or nitrogen supplements. This study has shown that it is possible to biologically treat Amarula distillery wastewater, a wastewater that has a high COD and high phenolic compound concentration, using white-rot fungus *Trametes pubescens* MB 89 and to obtain a high removal efficiency of COD and phenolic compounds (Strong, 2010).

Application of white-rot fungi in treatment of industrial effluents is summarized in Table 4.

Table 4. White-rot fungi used in decolorization of industrial effluents

White-rot fungi	Industry	References
<i>Coriolus versicolor</i>	textile	Hai et al., 2006
<i>Pleurotus ostreatus</i> , <i>Coriolus versicolor</i> and <i>Funalia trogii</i>	textile	Erkurt et al., 2007
<i>Coriolus versicolor</i>	textile	Asgher, 2009
<i>Pleurotus ostreatus</i>	textile	Lu et al., 2008
<i>Phanerochaete chrysosporium</i>	pulp and paper	Sharari et al., 2010
<i>Phanerochaete chrysosporium</i>	pulp and paper	Jaspers and Penninckx, 1996
* <i>Fomitopsis</i> sp. IMER2.	pulp and paper	Ma et al., 2008
<i>Phanerochaete chrysosporium</i> , <i>Pleurotus ostreatus</i> and S22	pulp and paper	Wu et al., 2005
<i>Pleurotus ostreatus</i>	food	Pant and Adholeya, 2007
<i>Pycnoporus coccineus</i>	food	Chairattananokorn et al., 2005
<i>Trametes pubescens</i> MB 89	food	Strong, 2010
<i>Pleurotus</i> sp.	food	Laconi et al., 2007
<i>Trametes versicolor</i> and <i>Pleurotus sajor</i>	food	Justino et al., 2009; Justino et al., 2010
<i>Panus tigrinus</i>	food	D'Annibale et al., 2006
<i>Phanerochaete chrysosporium</i>	coke	Lu et al., 2009

Conclusions

White-rot fungi possess complex and efficient lignolytic enzyme system. They have been successfully applied in treatment and decomposition of different phenolic compounds, dyes and other xenobiotics on the laboratory level. However, a lot of work is needed to be done to explore the potential of white-rot fungi in dye decolorization and for removal of hazardous chemicals on industrial scale. The paper underlines the importance of exploring potential of new strains in the process of bioremediation in order to expand the pool of

existing biocleaning and biobleaching white-rot fungi.

Acknowledgment

This research was financially supported by the Croatian Ministry of Science, Education and Sports (Contract Number 125-1252086-27939) and The National Foundation for Science, Higher Education and Technological Development of the Republic of Croatia.

References

- Acevedoa, F., Pizzul, L., Castillo, M.D.P., Cuevas, R., Diez, M.C. (2011): Degradation of polycyclic aromatic hydrocarbons by the Chilean white-rot fungus *Anthracophyllum discolor*, *J. Haz. Mat.* 185, 212–219.
- Aktaş, M., Çiçek, H., Taşpinaréal, A., Kibarér, G., Kolankoya, N., Tanyolaç, A. (2001): Reaction kinetics for laccase-catalyzed polymerization of 1-naphtol, *Biores. Technol.* 80, 29-36.
- Aktaş, M., Tanyolaç, A. (2003.a): Kinetics of laccase-catalyzed oxidative polymerization of catechol, *J. Mol. Catal. B.: Enzym.* 22, 61-69.
- Aktaş, M., Tanyolaç, A. (2003.b): Reaction conditions for laccase catalyzed polymerization of catechol, *Biores. Technol.* 87, 209-214.
- Alvarado, I.E., Navarro, D., Record, E., Asther, M., Aster, M., Lesaage-Meessen, L. (2003): Fungal biotransformation of *p*-coumaric acid into caffeic acid by *Pycnoporus cinnabarinus*: an alternative for producing a strong natural antioxidant, *Word J. Microbiol. Biotechnol.* 19, 157-160.
- Asgher, M., Bhatti, H.N., Ashraf, M., Legge, R.L. (2008): Recent developments in biodegradation of industrial pollutants by white-rot fungi and their enzyme system. *Biodegradation* 19, 771-783.
- Asgher, M., Azim, M., Bhatti, H.N. (2009): Decolorization of practical textile industry effluents by white-rot fungus *Coriolus versicolor* IBL-04, *Biochem. Eng. J.* 47, 61-65.
- Ander, P., Marzullo, L. (1997): Sugar oxidoreductases and veratryl alcohol oxidase as related to lignin degradation, *J. Biotechnol.* 53, 115-131.
- Bahri, H., Dignac, M.F., Rumpel, C., Rasse, D.P., Chenu, C., Mariotti, A. (2006): Lignin turnover kinetics in an agricultural soil is monomer specific; *Soil. Biol. Biochem.* 38, 1977-1988.
- Barclay, C.D., Legge R.L., Farquhar, G.F. (1993): Modelling the growth kinetics of *Phanerochaete chrysosporium* in submerged static cultures, *Appl. Environ. Microbiol.* 59, 1887-1892.
- Bending, G., Friloux, M., Walker, A. (2002): Degradation of contrasting pesticides by white-rot fungi and its relationship with ligninolytic potential, *FEMS Microbiol. Lett.* 18, 59-63.
- Bennet, J.W., Connick, W.J., Daigle, D., Wunch, K. (2001): Formulation of fungi for *in situ* bioremediation. In: *Fungi in Bioremediation*, Gadd, G.M. (ed.), New York, USA: Cambridge University Press, pp. 97-108.
- Bollag J.-M., Chu, H.-L., Rao, A., Gianfreda, L. (2003): Enzymatic oxidative transformation of chlorophenol mixtures. *J. Environ. Qual.* 32, 63-69.
- Boudet, A.-M. (2007): Evolution and current status of research in phenolic compounds, *Phytochemistry* 68, 2722-2735.
- Boyle, D., Wiesner, C., Richardson, A. (1998): Factors affecting the degradation of polycyclic aromatic hydrocarbons in soil by white-rot fungi, *Soil Biol. Biochem.* 30, 873–882.
- Burton, S.G. (2003): Laccases and phenol-oxidases in organic synthesis – a review, *Org. Chem.* 7, 1317-1331.
- Cerniglica, C.E., Sutherland, J.B. (2001): Bioremediation of polycyclic aromatic hydrocarbons by ligninolytic and non-ligninolytic fungi. In: *Fungi in Bioremediation*, Gadd, G.M. (ed.), New York, USA: Cambridge University Press, pp. 136-187.
- Chairattananokorn, P., Imai, T., Kondo, R., Sekine, M., Ukita, M. (2005): Decolorization of alcohol distillery wastewater by thermotolerant white-rot fungi, *Applied Biochem. Microbiol.* 41, 583–588.
- Chander, M., Arrora, D.S. (2007): Evaluation of some white-rot fungi for their potential to decolourise industrial dyes, *Dyes Pigments* 72, 192-198.
- Claus, H. (2003): Laccases and their occurrence in prokaryotes, *Arch. Microbiol.* 179, 145-150.
- Claus, H. (2004): Laccases: structure, reactions, distribution, *Micron.* 35, 93-96.
- Clemente, A.R., Anazawa, T.A., Durrant, L.R. (2001): Biodegradation of polycyclic aromatic hydrocarbons by soil fungi, *Braz. J. Microbiol.* 32, 255–261.
- Covert, S.F., Boldue, J., Cullen, D. (1992): Genomic organization of a cellulose gene family in *Phanerochaete chrysosporium*, *Curr. Genet.* 22, 407-413.
- D'Acunzo, F., Galli, C., Masci, B. (2002): Oxidation of phenols by laccase and laccase-mediator systems, *Eur. J. Biochem.* 269, 5330-5335.
- D'Annibale, A., Quaratino, D., Federici, F., Fenice, M. (2006): Effect of agitation and aeration on the reduction of pollutant load of olive mill wastewater by the white-rot fungus *Panus tigrinus*, *Biochem. Eng. J.* 29, 243–249.
- Dashtban, M., Schraft, H., Syed, T.A., Qin, W. (2010): Fungal biodegradation and enzymatic modification of lignin, *Int. J. Biochem. Mol. Biol.* 1, 36-50.
- De Wilde, C., Uzan, E., Zhou, Z., Kruus, K., Andberg, M., Buchert, J., Record, E., Asther, M., Lomascolo, A. (2008): Transgenic rice as a novel production system for *Melanocarpus* and *Pycnoporus* laccases, *Transgenic Res.* 17, 515–527.
- Eggen, T., Majcherczyk, A. (1998): Removal of polycyclic aromatic hydrocarbons (PAH) in contaminated soil by white-rot fungus *Pleurotus ostreatus*, *Int. Biodeg. Biodeg.* 41, 111-117.
- Eggen, T., Sveum, P. (1999): Decontamination of aged creosote polluted soil: the influence of temperature, white-rot fungus *Pleurotus ostreatus*, and pre-treatment, *Int. Biodeg. Biodeg.* 43, 125-133.
- Ergül, F.E., Sargin, S., Öngen, G., Sukan, F.V. (2010): Dephenolization and decolorization of olive mill wastewater through sequential batch and co-culture applications, *World J. Microbiol. Biotechnol.* DOI 10.1007/s11274-010-0433-4.

- Erkurt, E.A., Ünyayar, A., Kumbur, K. (2007): Decolorization of synthetic dyes by white-rot fungi, involving laccase enzyme in the process, *Proc. Biochem.* 42, 1429–1435.
- Evans, C.S., Hedger, J.N. (2001): Degradation of plant cell wall polymers. In: *Fungi in Bioremediation*, Gadd, G.M. (ed.), New York, USA: Cambridge University Press, pp. 1-26.
- Fang, Z., Sato, T., Smith, Jr.R.L., Inomata, H., Arai, K., Kozinski, J.A. (2008): Reaction chemistry and phase behavior of lignin in high-temperature and supercritical water, *Bioresour. Technol.* 99, 3424–3430.
- Fan, F., Zhuo, R., Sun, S., Wan, X., Jiang, M., Zhang, X., Yang, Y. (2010): Cloning and functional analysis of a new laccase gene from *Trametes* sp. 48424 which had the high yield of laccase and strong ability for decolorizing different dyes, *Bioresour. Technol.*, DOI 10.1016/j.biortech.2010.10.079.
- Faraco, V., Pezzella, C., Miele, A., Giardina, P., Sannia, G. (2009): Bio-remediation of colored industrial wastewaters by the white-rot fungi *Phanerochaete chrysosporium* and *Pleurotus ostreatus* and their enzymes, *Biodegradation*, 20, 209–220.
- Gao, D., Du, L., Yang, J., Wu, W-M., Liang, H. (2010): A critical review of the application of white-rot fungus to environmental pollution control, *Crit. Rev. Biotechnol.* 30, 70–77.
- Geng, X., Li, K. (2002): Degradation of non-phenolic lignin by the white-rot fungus *Pycnoporus cinnabarinus*, *Appl. Microbiol. Biotechnol.* 60, 342–346.
- Gianfreda, L., Xum F., Bollag, J.M. (1999): Laccase - A useful group of oxidoreductive enzymes, *Biorem. J.* 3, 1-25.
- Gianfreda, L., Sannino, F., Rao, M.A., Bolla, J.-M. (2003): Oxidative transformation of phenols in aqueous mixture, *Water Res.* 37, 3205-3215.
- Gianfreda, L., Iamarino, G., Scelza, R., Rao, M.A. (2006): Oxidative catalysts for the transformation of phenolic pollutants: a brief review, *Biocatal. Biotrans.* 24, 177-187.
- Gomes Machado, K.M., Matheus, D.R., Rosim Monteiro, R.T., Ramos Bononi, V.L. (2005): Biodegradation of pentachlorophenol by tropical basidiomycetes in soils contaminated with industrial residues, *World J. Microb. Biotechnol.* 21, 297–301.
- Grönqvist, S., Viikari, L., Niku-Paavola, M.-L., Orlandi, M., Canevali, C., Buchert, J. (2005): Oxidation of milled wood lignin with laccase, tyrosinase and horseradish peroxidase, *Appl. Microbiol. Biotechnol.* 67, 489-494.
- Gupta, R., Mehta, G., Khasa, Y.P. (2010): Fungal delignification of lignocellulosic biomass improves the saccharification of cellulotics, *Biodegradation*, DOI 10.1007/s10532-010-9404-6.
- Güreşir, M., Aktaş, N., Tanyolaç, A. (2005): Influence of reaction conditions on the rate of enzymatic polymerization of pyrogallol using laccase, *Process Biochem.* 40, 1175-1182.
- Hai, F.I., Yamamoto, K., Fukushi, K. (2006): Development of a submerged membrane fungi reactor for textile wastewater treatment, *Desalination* 192, 315–322.
- Hamid, M., Rehman, K. (2009): Potential applications of peroxidases, *Food Chem.* 115, 1177–1186.
- Harvey, P., Thurston, C.F. (2001): The biochemistry of lignolytic fungi. In: *Fungi in Bioremediation*, Gadd, G.M. (ed.), New York, USA: Cambridge University Press, pp. 27-51.
- Hatakka, A. (1994): Lignin-modifying enzymes from selected white-rot fungi: production and role in lignin degradation, *FEMS Microbiol. Rev.* 13, 125-135.
- Haritash, A.K., Kaushik, C.P. (2009): Biodegradation aspects of polycyclic aromatic hydrocarbons (PAHs): A review, *J. Haz. Mat.* 169, 1-15.
- Higson, F.K. (1991): Degradation of xenobiotics by white-rot fungi, *Rev. Environ. Contam. Toxicol.* 122, 111–152.
- Hofrichter, M. (2002): Review: lignin conversion by manganese peroxidase (MnP), *Enzyme Microb. Technol.* 30, 454-466.
- Jang M.Y., Ryu W.R., Cho, M.H. (2002): Laccase production from repeated batch culture using free mycelia of *Trametes* sp., *Enzyme. Microb. Technol.* 33, 741-746.
- Jaouani, A., Sayadi S., Vanthourhout, V., Penninckx, M.J. (2003): Potent fungi for decolourisation of olive oil mill wastewaters, *Enzyme. Microb. Technol.* 33, 802-809.
- Jaspers, C.J., Penninckx, M.J. (1996): Adsorption effects in the decolorization of a kraft bleach plant effluent by *Phanerochaete chrysosporium*, *Biotechnol. Lett.* 18, 1257-1260.
- Jonas, U., Hammer, E., Haupt, E. T. K., Schauer, F. (2000): Characterization of coupling products formed by biotransformation of biphenyl and diphenyl ether by the white-rot fungus *Pycnoporus cinnabarinus*, *Arch. Microbiol.* 174, 393–398.
- Justino, C., Marques, A.G., Duarte, K.R., Costa Duarte, A., Pereira, R., Rocha-Santos, T., Freitas, A.C. (2010): Degradation of phenols in olive oil mill wastewater by biological, enzymatic, and photo-Fenton oxidation, *Environ. Sci. Pollut. Res.* 17, 650-656.
- Justino, C.I., Duarte, K., Loureiro, F., Pereira, R., Antunes, S.C., Marques, S.M., Gonçalves, F., Rocha-Santosa, T.A.P., Freitas, A.C. (2009): Toxicity and organic content characterization of olive oil mill wastewater undergoing a sequential treatment with fungi and photo-Fenton oxidation, *J. Hazard. Mater.* 172, 1560-1572.
- Kariminiaae-Hamedani, H., Sakurai, A., Sakakibara, M. (2007): Decolorization of synthetic dyes by a new manganese peroxidase-producing white-rot fungus, *Dyes Pigments* 72, 157-162.

- Kasai, N., Ikushiro, S., Shinkyō, R., Yasuda, K., Hirose, S., Arisawa, A., Ichinose, H., Wariishi, H., Sakaki, T. (2010): Metabolism of mono- and dichloro-dibenzo-*p*-dioxins by *Phanerochaete chrysosporium* cytochromes P450, *Appl. Microbiol. Biotechnol.* 86, 773-780.
- Klaus, P. (2002): New insights into lignin peroxidase, *Ind. J. Chem.* 41, 46-53.
- Knapp, J.S., Vantoch-Wood, E.J., Zhang, F. (2001): Use of wood-rotting fungi for the decolorization of dyes and industrial effluents. In: *Fungi in bioremediation*, Gadd, G.M. (ed.), New York, USA: Cambridge University Press, pp. 242-304.
- Kurniawati, S., Nicell J.A. (2006): A pseudo-steady state model of the kinetics of laccase-catalysed oxidation of aqueous phenol, *J. Chem. Technol. Biotechnol.* 81, 1198-1208.
- Kurniawati, S., Nicell J.A. (2007): Efficacy of mediators for enhancing the laccase-catalyzed oxidation of aqueous phenol, *Enzyme Microb. Technol.* 41, 353-361.
- Kurniawati, S., Nicell J.A. (2008): Characterization of *Trametes versicolor* laccase for the transformation of aqueous phenol, *Biores. Technol.* 99, 7825-7834.
- Laconi, S., Molle, G., Cabiddu, A., Pompei, R. (2007): Bioremediation of olive oil mill wastewater and production of microbial biomass, *Biodegradation* 18, 559-566.
- Lara, M.A., Rodríguez-Malaver, J., Rojas, O.J., Hoimquist, O., González, A.M., Bullón, J., Peñaloza, N., Araujo, E. (2002): Black liquor lignin biodegradation by *Trametes elegans*, *Int. Biodeter. Biodegr.* 52, 167-173.
- Leisola, M.S.A., Garcia, S. (1989): The mechanism of lignin degradation. In: *Enzyme Systems for Lignocellulose Degradation*, Coughlan, M.P. (ed.), New York, USA, Elsevier Science Publishing co., inc., pp. 89-109
- Leisola, M., Ulmer, D., Fiechter, A. (1984): Variations in some extracellular enzyme activities during degradation of lignocellulose by *Phanerochaete chrysosporium*, *Appl. Biochem. Biotechnol.* 9, 373-374.
- Levin, L., Herrmann, C., Papinutti, V.L. (2008): Optimization of lignocellulolytic enzyme production by the white-rot fungus *Trametes trogii* in solid-state fermentation using response surface methodology, *Biochem. Eng. J.* 39, 207-214.
- Levin, L., Malignani, E., Ramos, A.M. (2010): Effect of nitrogen sources and vitamins on ligninolytic enzyme production by some white-rot fungi. Dye decolorization by selected culture filtrates, *Bioresour. Technol.* 101, 4554-4563.
- Linko, S. (1992): Production of *Phanerochaete chrysosporium* lignin peroxidase, *Biotechnol. Adv.* 10, 191-236.
- López, C., Mielgo, I., Moreira, M.T., Feijoo, G., Lema J.M. (2002): Enzymatic membrane reactors for biodegradation of recalcitrant compounds. Application to dye decolorization, *J. Biotechnol.* 99, 249-257.
- Lu, Y., Yan, L., Wang, Y., Zhou, S., Fu, J., Zhang, J. (2009): Biodegradation of phenolic compounds from coking wastewater by immobilized white-rot fungus *Phanerochaete chrysosporium*, *J. Haz. Mater.* 165, 1091-1097.
- Lu, Y., Phillips, D.R., Lu, L., Hardin, I.R. (2008): Determination of the degradation products of selected sulfonated phenylazonaphthol dyes treated by white-rot fungus *Pleurotus ostreatus* by capillary electrophoresis coupled with electrospray ionization ion trap mass spectrometry, *J. Chromatogr. A* 1208, 223-231.
- Lucas, M., Mertens, V., Corbisier, A.-M., Vanhulle, S. (2007): Synthetic dyes decolourisation by white-rot fungi: Development of original microtitre plate method and screening, *Enzyme Microb. Technol.* 42, 97-106.
- Lundell, T.K., Mäkelä, M.R., Hildén, K. (2010): Lignin-modifying enzymes in filamentous basidiomycetes – ecological, functional and phylogenetic review, *J. Basic Microbiol.* 50, 5-20.
- Ma, F., Xiong, Z., Zheng, Y., Yu, X., Zhang, X. (2008): Repeated batch process for biological treatment of black liquor using brown-rot basidiomycete *Fomitopsis* sp. IMER2, *World J. Microbiol. Biotechnol.* 24, 2627-2632.
- Maloney, S.E. (2001): Pesticide degradation. In: *Fungi in bioremediation*, Gadd, G.M. (ed.), New York, USA: Cambridge University Press, 188-223.
- Marco-Urrea, E., Pérez-Trujillo, M., Vicent, T., Caminal, G. (2009): Ability of white-rot fungi to remove selected pharmaceuticals and identification of degradation products of ibuprofen by *Trametes versicolor*, *Chemosphere*, 74, 765-772.
- Mayer, A.M., Staples, R.C. (2002): Laccase: new function for an old enzyme, *Phytochemistry* 60, 551-55.
- Membrillo, I., Sánchez, C., Meneses, M., Favela, E., Loera, O. (2008): Effect of substrate particle size and additional nitrogen source on production of lignocellulolytic enzymes by *Pleurotus ostreatus* strains, *Biores. Technol.* 99, 7842-7847.
- Moreira, M.T., Sierra-Alvarez, R., Lema, J.M., Feijoo, G., Field, J.A. (2001): Oxidation of lignin in eucalyptus kraft pulp by manganese peroxidase from *Bjerkandera* sp. Strain BOS55, *Biores. Technol.* 78, 71-79.
- More, T.T., Yan, S., Tyagi, R.D., Surampalli, R.Y. (2010): Potential use of filamentous fungi for wastewater sludge treatment, *Biores. Technol.* 101, 7691-7700.
- Mussatto, S.I., Fernandes, M., Roberto, I.C. (2007): Lignin recovery from brewer's spent black liquor, *Carbohydr. Polym.* 70, 218-223.
- Nilsson, I., Möller, A., Mattiasson, B., Rubindamayugi, M.S.T., Welander, U. (2006): Decolorization of synthetic and real textile wastewater by the use of white-rot fungi, *Enzyme Microb. Technol.* 38, 94-100.
- Palmieri, G., Giardina, P., Bianco, C., Scaloni, A., Capasso, A., Sannia, G. (1997): A novel white laccase from *Pleurotus ostreatus*, *J. Biol. Chem.* 272, 31301-31307.

- Pant, D., Adholeya, A. (2007): Enhanced production of ligninolytic enzymes and decolorization of molasses distillery wastewater by fungi under solid state fermentation, *Biodegradation* 18, 647-659.
- Papagianni, M. (2004): Fungal morphology and metabolite production in submerged mycelial processes, *Biotechnol. Adv.* 22, 189-259.
- Podgornik, H., Poljanšek, I., Perdih, A. (2001a): Transformation of Indigo carmine by *Phanerochaete chrysosporium* ligninolytic enzyme, *Enzyme Microb. Technol.* 29, 166-172.
- Podgornik, H., Stegu, M., Podgornik, A., Perdih, A. (2001b): Isolation and characterization of Mn(III) tartarate from *Phanerochaete chrysosporium* culture broth, *FEMS Microbiol. Lett.* 201, 265-269.
- Podgornik, H., Podgornik, A. (2002): Characteristics of LiP immobilized to CIM monolithic supports, *Enzyme Microb. Technol.* 31, 855-861.
- Quintero, J.C., Lú-Chau, T.A., Moreira, M.T., Feijoo, G., Lema, J. (2007): Bioremediation of HCH present in soil by the white-rot fungus *Bjerkandera adusta* in a slurry batch bioreactor, *Int. Biodet. Biodeg.* 60, 319-326.
- Rancano, G., Lorenzo, M., Molares, N., Rodríguez Couto, S., Sanromán, A. (2003): Production of laccase by *Trametes versicolor* in an airlift fermentor, *Proc. Biochem.* 39, 467-473.
- Reddy, G.V., Ravindra Babu, P., Komaraiah, P., Roy, K.R.R.M., Kothari, I.L.: Utilization of banana waste for the production of lignolytic and cellulolytic enzymes by solid substrate fermentation using two *Pleurotus* species (*P. ostreatus* and *P. sajor-caju*), *Process Biochem.* 38, 1457-1462.
- Renganathan, V., Usha, S.N., Lindenburg, F. (1990): Cellobiose-oxidizing enzymes from the lignocellulose-degrading basidiomycete *Phanerochaete chrysosporium*: interaction with microcrystalline cellulose, *Appl. Microbiol. Biotechnol.* 32, 609-613.
- Revankar, M.S., Lele, S.S. (2006): Increased production of extracellular laccase by the white-rot fungus *Coriolus versicolor* MTCC 138, *World J. Microbiol. Biotechnol.* 22, 921-926.
- Rivela, I., Rodríguez Couto, S., Sanromán, A. (2000): Extracellular ligninolytic enzyme production by *Phanerochaete chrysosporium* in a new solid-state bioreactor, *Biotechnol. Lett.* 22, 1443-1447.
- Rodríguez Couto, S., Feijoo, G., Moreira, M.T., Lema, J.M. (2002): Evaluation of the environmental conditions for the continuous production of lignin peroxidase by *Phanerochaete chrysosporium* in fixed-bed bioreactors, *Biotechnol. Lett.* 24, 791-794.
- Rodríguez Couto, S., Toca-Herrera, J.L. (2006): Industrial and biotechnological application of laccase: A review, *Biotechnol. Adv.* 24, 500-513.
- Rodríguez Couto, S., Toca-Herrera, J.L. (2007): Laccase production at reactor scale by filamentous fungi, *Biotechnol. Adv.* 25, 558-569.
- Rodríguez Couto, S., Gundín, M., Lorenzo, M., Sanromán, M.A. (2007): Screening of supports and inducers for laccase production by *Trametes versicolor* in semi-solid-state conditions, *Proc. Biochem.* 38, 249-255.
- Rodríguez-Rodríguez, C.E., Marco-Urrea, E., Caminal, G. (2010): Naproxen degradation test to monitor *Trametes versicolor* activity in solid-state bioremediation processes, *J. Haz. Mat.* 179, 1152-1155.
- Singh, H. (2006): Mycoremediation, Fungal Bioremediation, New Jersey, USA: John Wiley and Sons, pp. 1-28, 29-75.
- Singh, S., Pakshirajan, K., Daverey, A. (2009): Enhanced decolorization of Direct Red-80 dye by the white-rot fungus *Phanerochaete chrysosporium* employing sequential design of experiments, *Biodegradation* 21, 501-511.
- Singh, S., Pakshirajan, K. (2010): Enzyme activities and decolorization of single and mixed azo dyes by the white-rot fungus *Phanerochaete chrysosporium*, *Int. Biodet. Biodeg.* 64, 146-150.
- Sigoillot, J.C., Herpoël, I., Frasse, P., Moukha, S., Lesage-Meessen, L., Marcel, A. (1999): Laccase production by a monokaryotic strain of *Pycnoporus cinnabarinus* derived from a dikaryotic strain, *World J. Microbiol. Biotechnol.* 15, 481-484.
- Shahvali, M., Assadi, M.M., Rostani, K. (2000): Effect on environmental parameters on decolorization of textile wastewater using *Phanerochaete chrysosporium*, *Bioproc. Eng.* 23, 721-726.
- Sharari, M., Jahan Latibari, A., Guillet, A., Aourousea, M., Mouhamadou, B., Rafeiee, G., Mirshokraei, A., Parsapaghough, D. (2010): Application of the white-rot fungus *Phanerochaete chrysosporium* in biotreatment of bagasse effluent, *Biodegradation*, DOI 10.1007/s10532-010-9415-3
- Strong, P. J. (2010): Fungal remediation of Amarula distillery wastewater, *World J. Microbiol. Biotechnol.* 26, 133-144.
- Tagger, S., Périssol, C., Gil, G., Vogt, G., Petit J.L. (1998): Phenoloxidase of the white-rot fungus *Marasmius quercophilus* isolated from an evergreen oak litter (*Quercus ilex* L.), *Enzyme Microb. Technol.* 23, 372-379.
- Takano, M., Abe, H., Hayashi, N. (2006): Extracellular peroxidase activity at the hyphal tips of the white-rot fungus *Phanerochaete crassa* WD1694, *J. Wood Sci.* 52, 429-435.
- Tavares, A.P.M., Coelho, M.A.Z., Coutinho, J.A.P., Xavier, A.M.R.B. (2005): Laccase improvement in submerged cultivation induced production and kinetic modeling, *J. Chem. Technol. Biotechnol.* 80, 669-676.
- Teerapatsakul, C., Parra, R., Bucke, C., Chitradon, L. (2007): Improvement of laccase production from *Ganoderma* sp. KU-alk4 by medium engineering, *World J. Microbiol. Biotechnol.* 23, 1519-1527.
- Thiruchelvam, A.T., Ramsay, A.J. (2007): Growth and laccase production kinetics of *Trametes versicolor* in stirred tank reactor, *Appl. Microbiol. Biotechnol.* 74, 547-554.

- Tišma, M., Žnidaršič-Plazl, P., Plazl, I., Zelić, B., Vasić-Rački, Đ. (2008): Modelling of L-DOPA oxidation catalyzed by laccase, *Chem. Biochem. Eng. Q.* 22, 134-142.
- Tišma, M., Zelić, B., Vasić-Rački, Đ., Žnidaršič-Plazl, P., Plazl, I. (2009): Modelling of laccase-catalyzed L-DOPA oxidation in a microreactor, *Chem. Eng. J.* 149, 383-388.
- Tišma, M., Sudar, M., Vasić-Rački, Đ., Zelić, B. (2010a): Mathematical model for *Trametes versicolor* growth in submerged cultivation, *Bioproc. Biosyst. Eng.* 33, 749-758.
- Tišma, M., Zelić, B., Vasić-Rački, Đ., Žnidaršič-Plazl, P., Plazl, I. (2010b): Oxidation of coniferyl alcohol catalyzed by laccases from *Trametes versicolor*, *Acta Chim. Slo.* 57, 110-117.
- Tortella, G.R., Rubilar, O., Gianfreda, L., Valenzuela, E., Diez, M.C. (2008): Enzymatic characterization of Chilean native wood-rotting fungi for potential use in the bioremediation of polluted environments with chlorophenols, *World J. Microbiol. Biotechnol.* 24, 2805-2818.
- Tychanowicz, G. K., Zilly, A., de Souza, C.G.M., Peralta, R.M. (2004): Decolorization of industrial dyes by solid-state cultures of *Pleurotis pulmonarius*, *Process Biochem.* 39, 855-859.
- Valentín, L., Feijoo, G., Moreira, M.T., Lema, J.M. (2006): Biodegradation of polycyclic aromatic hydrocarbons in forest and salt marsh soils by white-rot fungi, 58, 15-21.
- Valentín, L., Lu-Chau, T.A., López, C., Feijoo, G., Moreira, M.T., Lema, J.M. (2007): Biodegradation of dibenzothiophene, fluoranthene, pyrene and chrysene in a soil slurry reactor by the white-rot fungus *Bjerkandera* sp. BOS55, *Process Biochem.* 42, 641-648.
- Vijaykumar, M.H., Veeranagouda, Y., Neelakanteshwar, K., Karegoudar, T.B. (2006): Decolorization of 1:2 metal complex dye Acid blue 193 by a newly isolated fungus *Cladosporium cladosporioides*, *World J. Microbiol. Biotechnol.* 22, 157-162.
- Wang, H., Lu, F., Sun, Y., Du, L. (2004): Heterologous expression of lignin peroxidase of *Phanerochaete chrysosporium* in *Pichia methanolica*, *Biotechnol. Lett.* 26, 1569-1573.
- Wang, C., Xi, J-Y, Hu, H-Y, Wen, X-H. (2008): Biodegradation of gaseous chlorobenzene by white-rot fungus *Phanerochaete chrysosporium*, *Biomed. Environ. Sci.* 21, 474-478.
- Webster, J., Weber, R.W.S. (2007): Introduction to Fungi, New York, USA: Cambridge University Press, pp. 1.
- Wesenberg, D. Kyriakides, I., Agathos, S. N. (2003): White-rot fungi and their enzymes for the treatment of industrial dye effluents, *Biotechnol. Adv.* 22, 161-187.
- Widsten, P., Kandelbauer, A. (2008): Laccase applications in the forest products industry: A review, *Enzyme Microb. Technol.* 42, 293-307.
- Wu, J., Xiao, Y.Z., Yu, H.Q. (2005): Degradation of lignin in pulp mill wastewaters by white-rot fungi on biofilm, *Biores. Technol.* 96, 1357-1363.
- Xavier, A.M.R.B., Tavares A.P.M., Ferreira, R., Amado, F. (2007): *Trametes versicolor* growth and laccase induction with by-products of pulp and paper industry, *E. J. Biotechnol.* 10, 444-451.
- Zille, A., Munteanu, F.D., Gübitz, G.M., Cavaco-Paulo, A. (2005): Laccase kinetics of degradation and coupling reaction, *J. Mol. Catal. B: Enzymatic* 33, 23-28.
- Zhang, F., Knapp, J.S., Tapley, K.N. (1998): Decolourisation of cotton bleaching effluent in a continuous fluidized-bed bioreactor using wood rotting fungus, *Biotechnol. Lett.* 20, 717-723.
- Žnidaršič, P., Pavko A. (2001): The morphology of filamentous fungi in submerged cultivation as a bioprocess parameter, *Food Technol. Biotechnol.* 39, 237-252.
- Žnidaršič-Plazl, P. (2006): The influence of some engineering variables upon the morphology of *Rhizopus nigricans* in a stirred tank bioreactor, *Chem. Biochem. Eng. Q.* 20, 275-280.

Received: November 03, 2010

Accepted: December 10, 2010