

Time-course experiment of Fusarium infestation of wheat genotypes with the emphasis on the physiological response

Marček, Tihana; Viljevac Vuletić, Marija; Bakula, Ivan; Alivojvodić, Sara; Španić, Valentina

Source / Izvornik: **Croatian journal of food science and technology, 2018, 10, 58 - 63**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.17508/CJFST.2018.10.1.11>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:109:200466>

Rights / Prava: [Attribution-NoDerivatives 4.0 International](#)/[Imenovanje-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-02-23**

REPOZITORIJ

PTFS

PREHRAMBENO-TEHNOLOŠKI FAKULTET OSIJEK

dabar
DIGITALNI AKADEMSKI ARHIVI I REPOZITORIJI

Repository / Repozitorij:

[Repository of the Faculty of Food Technology Osijek](#)





Original scientific paper

DOI: 10.17508/CJFST.2018.10.1.11

Time-course experiment of *Fusarium* infestation of wheat genotypes with the emphasis on the physiological response

TIHANA MARČEK^{1*}, MARIJA VILJEVAC VULETIĆ², IVAN BAKULA¹, SARA ALIVOJVODIĆ¹, VALENTINA ŠPANIĆ²

¹Josip Juraj Strossmayer University of Osijek, Faculty of Food Technology Osijek, Department of Food and Nutrition Research, F. Kuhača 20, 31000 Osijek, Croatia

²Agricultural Institute Osijek, Južno predgrađe 17, 31000 Osijek, Croatia

ARTICLE INFO

Article history:

Received: January 25, 2018

Accepted: February 24, 2018

Keywords:

wheat,
Fusarium spp.,
MDA,
H₂O₂,
antioxidative enzymes.

ABSTRACT

Wheat (*Triticum aestivum* L.) is one of the most significant food cereal in the world. Under natural conditions, biotic and abiotic stress factors can seriously endanger the plant growth and development. Fusarium head blight (FHB), mainly caused by *F. graminearum* and *F. culmorum*, is a disease that has negative effects on economy, namely on the yield and the quality of the grain. In this research, the activities of guaiacol peroxidase (POD), catalase (CAT), ascorbate peroxidase (APX) and polyphenol oxidase (PPO), as well as H₂O₂ concentration and malondialdehyde (MDA) content, were determined in three wheat genotypes of various resistance to FHB ("Super Žitarka"-sensitive; "Lucija"- moderately resistant, and "Apache"- resistant) after the 2nd and the 4th day of infestation. The infected plants of "Lucija" showed higher POD activity compared to control after the 4th day, and increased POD activity compared to the 2nd day of infection. On the 2nd day, "Apache" responded to pathogen by the enhanced APX activity, and higher MDA level, compared to the 4th day of treatment when it started to decrease. Moreover, the treated plants of the same genotype showed the increased MDA level on the 2nd day, unlike the untreated plants. Contrarily, H₂O₂ concentration in "Apache" showed the tendency to increase with the time of *Fusarium*-exposure. In comparison to the 2nd day of treatment, PPO activity decreased in "Super Žitarka" on the 4th day. In all genotypes at both sampling points, CAT activity did not change significantly under the pathogen attack. Overproduction of H₂O₂ accompanied with the increased APX activity 2 days after infection may lead to the conclusion that FHB tolerance of "Apache" is ensured by the earlier activation of the defence mechanisms.

Introduction

Fusarium head blight (FHB) caused by *Fusarium* species is an economically important wheat (*Triticum aestivum* L.) disease that has a negative impact on the yield and grain quality. Moderate to warm temperatures in conjunction with high humidity during the stage of anthesis and early kernel development increase the possibility of infection (Gilbert and Tekauz, 2000). Infected kernels become bleached, while contaminated grains

are often shrivelled with the presence of mycotoxins that have a harmful effect on human and animal health (Bai and Shaner, 2004).

In nature, plant copes with a numerous biotic and abiotic stresses. Biotic stresses include plant infestation by different pathogen elicitors or insects. Exposure of plants to *Fusarium* infection results in the activation of plant defence mechanisms (Nanda et al., 2010). These mechanisms can provide the efficient protection for the plant against reactive oxygen species (ROS) in the manner they remove them. ROS pool consists of H₂O₂, singlet oxygen (¹O₂), superoxide (O₂⁻) and hydroxyl (OH[•]) radicals

*Corresponding author E-mail:
tihana.marcek@ptfos.hr

and their generation in plant cell depends on the environmental conditions (Thannickal and Fanburg, 2000). Overaccumulation of ROS in plant leads to a disruption of the membranes' phospholipid bilayer, DNA damages and protein denaturation, thereby affecting the physiological processes in plants (Dat et al., 2000; García-Limones et al., 2002). Under the stress exposure, plant activates an enzymatic antioxidative system, which includes catalase (CAT), ascorbate peroxidase (APX) and non-specific peroxidase (POD), as well as nonenzymatic antioxidative detoxifying enzymes such as polyphenol oxidase (PPO). Major role of PPO is to eliminate ROS using flavonoids and phenolics as substrates (Boeckx et al., 2015; Mayer, 2006). There is a growing body of evidence suggesting that high induction of antioxidative enzymes in plants contributes to the pathogen resistance (Madadkhah et al., 2012; Racchi, 2013; Shahbazi et al., 2010; Sorahinobar et al., 2015). H₂O₂ has the poorest reactivity and ability to diffuse easily across the membranes, which distinguishes it from other ROS particles. One of the roles of H₂O₂ under pathogen attack is to reduce the growth and viability of pathogen in order to restrict its entrance into plant (Kuzniak and Urbanek, 2000).

The aim of this research was to explore the effect of *Fusarium* spp. in three wheat genotypes with different FHB resistance ("Super Žitarka"-sensitive; "Lucija"- moderately resistant, and "Apache"- resistant) after the 2nd and the 4th day of infestation by determining the activity of the antioxidative enzymes (APX, CAT, and POD), lipid peroxidation content, H₂O₂ level and PPO activity.

Materials and methods

Field trials and pathogen inoculum preparation

The experiment was carried out under field conditions. Wheat spikes of the wheat genotypes ("Super Žitarka", "Apache" and "Lucija") were treated with a mixture of *Fusarium graminearum* and *Fusarium culmorum* (stressed plants), while the second group of plants were left to natural infection (control). *Fusarium* treatment was performed by spraying the spikes at flowering (Zadok's scale 65) (Zadoks et al., 1974) using a tractor back sprinkler in the afternoon hours. The plants were irrigated few times during the day in order to maintain moisture on the ears.

After the 2nd and 4th day of *Fusarium* treatment, spikes were collected and stored at -80 °C prior to analyses. For the antioxidative enzymes activity, fresh tissue (200 mg) was ground into fine powder by using pestle and mortar, homogenized in

50 mmol/dm³ potassium phosphate buffer (pH 7.0) by adding 0.1 mmol/dm³ of ethylenediaminetetraacetic acid, 5 mmol/dm³ of ascorbate acid and polyvinylpolypyrrolidone (PVP), centrifuged at 14 000 rcf for 30 min at 4 °C. From the obtained supernatants, the enzyme activity and the total protein content (Bradford, 1976) were measured. The activity of enzymes was expressed as units (U) per milligram of proteins [U mg⁻¹ proteins].

Activity of antioxidative enzymes

The POD activity was assayed at 470 nm following the method by Siegel and Galson (1967) as peroxidation of hydrogen peroxide with guaiacol as an electron donor. CAT activity was determined by the decline in absorbance at 240 nm (Aebi, 1984). APX activity was analysed by tracking the decline in absorbance of ascorbate at 290 nm (Nakano and Asada, 1981). PPO activity was estimated at 430 nm as a rate of oxidation of pyrogallol by the increase in absorbance (Raymond et al., 1993). Enzymes' activity was presented as units (U) of enzyme activity per milligram of protein [μmol min/mg/proteins].

MDA content and H₂O₂ concentration

For the MDA and H₂O₂ determination, the spikes (400 mg) were homogenized in 0.1% (w/v) trichloroacetic acid (TCA) and centrifuged at 12 000 rcf for 15 min at 4 °C. The MDA content, final product of the lipid peroxidation, was determined according to protocol by Verma and Dubey (2003).

5 % TBA in 20% TCA was added to 0.5 mL of supernatant, heated at 95 °C for 30 min and cooled in an ice bath. After centrifugation (14 000 rcf for 15 min at 4°C), absorbance of supernatant was measured at 532 nm and 600 nm. The data for non-specific absorption at 600 nm was subtracted. As a blank, the 0.5 % TBA in 20 % TCA solution was used. The MDA content was calculated using the molar extinction coefficient as 155 mmol/dm³/cm.

H₂O₂ concentration was determined using the protocol by Velikova et al. (2000). The 0.5 mL of extract was added to the mixture of 10 mmol/dm³ potassium phosphate buffer (pH 7.0) and 1 mol/dm³ KI and kept in darkness for 20 min. Absorbance of the mixture was read at 390 nm. H₂O₂ content was defined using a calibration curve obtained with H₂O₂ solutions ranging from 20 to 700 μmol/dm³ and expressed as millimoles per gram of fresh weight [mmol/g_{FW}].

All results were expressed as means of five replicates with the corresponding standard errors. Differences between means were evaluated using the nonparametric Kruskal-Wallis test using Statistica 13 software.

Results and discussion

In plants, increased POD and PPO activity are mostly related to the straightening of the cell wall in order to restrict the pathogen entrance (Mohammadi and Karr, 2002; Morkunas and Gemerek, 2007). In stressed plants of “Apache”, POD activity increased over time (it was higher on the 4th day in comparison to the 2nd day of infection) (Fig. 1A). Such a delayed reaction could be the result of the stronger macroconidia colonization and penetration of the host plant cells. Similar result was observed in FHB-resistant wheat genotype “Vulkan”, in which *Fusarium* treated plant exhibited higher POD induction at a later point of collection (96 hai) than at 48 hai (Španić et al., 2017). The partially FHB-resistant genotype “Lucija” showed an increase in POD activity under *Fusarium* attack on the 4th day in regard to respective control group (Fig. 1A) suggesting that POD in “Lucija” is involved in overcoming stress injuries provoked by pathogen. The increase of POD activity after the 4th day of

infection was also described in tomato leaves exposed to *Botrytis cinerea* (Kuźniak and Skłodowska, 2005).

Apart from the involvement in ROS removal, the role of PPO is connected with non-enzymatic plant defence response against pathogen (Boeckx et al., 2015; Parveen et al., 2010; Thipyapong et al., 2004). In all tested genotypes, PPO activity did not show significant changes during time-course experiment (Fig. 1B). Exception was FHB-sensitive “Super Žitarka”, which responded to pathogen elicitor by an enhanced expression of PPO activity at an earlier time-point (2nd day) compared to later sampling time (4th day), which lead us to conclusion that “Super Žitarka” activates the non-enzymatic antioxidative system in response reaction earlier. On the other hand, FHB-sensitive wheat cultivar “Falat” showed induction of PPO activity after 5 days of *Fusarium* stress (Sorahinobar et al., 2015). The same was noticed in roots of *Fusarium*-susceptible melon genotypes on the 4th day of exposure to *F. oxysporum* f. sp. *melonis* (Madadkhah et al., 2012).




Fig. 1. Activity of POD (A), PPO (B) APX (C) and CAT (D) in ears of wheat genotypes (“Super Žitarka”, “Apache” and “Lucija”) infected with *Fusarium* spp. after 2nd and 4th day (C-control, T-treatment). Values are means of five replicates \pm SE. Different capital letters indicate significantly different values ($p < 0.05$) among different time points and different genotypes under the same treatment, while different lower-case letters represent significantly different values ($p < 0.05$) within each genotype separately at same sampling point.




Fig. 2. H₂O₂ (A) and MDA content (B) in ears of wheat genotypes “Super Žitarka”, “Apache” and “Lucija” infected with *Fusarium* spp. after 2nd and 4th day (C-control, T-treatment). Values are means of five replicates ±SE. Different capital letters indicate significantly different values ($P < 0.05$) among different time points and different genotypes under the same treatment, while different lower-case letters represent significantly different values ($P < 0.05$) within each genotype separately at same sampling point.

In “Apache”, at an early time point (2nd day) of infection, activity of APX was higher than on the 4th day (Fig. 1C) indicating that this genotype involved APX more in early response to *Fusarium*. The rapid APX activity under *Fusarium* treatment was also noticed in FHB-resistant wheat genotype “Vulkan” (Španić et al., 2017). Moreover, in “Apache”, H₂O₂ concentration (Fig. 2A) was lower on the 2nd day of infection compared to the 4th day, which could be the result of an increased APX activity suggesting that APX had an important role in the early ROS decomposition. The same was found in pea (*Pisum sativum* L.) - *Sclerotinia sclerotiorum* interaction (Jain et al., 2013). The great importance of APX in ensuring the disease resistance was also noticed in *Avena sativa* subjected to *Puccinia coronata* stress (Figueiró et al., 2015).

Considering H₂O₂ production under stressful environment, “Apache” exhibited overproduction of H₂O₂ on the 4th day in comparison to the 2nd day (Fig. 2A). Higher H₂O₂ accumulation was found 72 h after the infection of *Fusarium* tolerant-iso banana genotype with *Fusarium oxysporum* (Li et al., 2011). In our study, overproduction of H₂O₂ in “Apache” under pathogen stress could be the result of its participation in plant defence mechanisms. Recent reports indicate that in response to elicitor, plant can increase H₂O₂ level in order to induce the defence (Vranova et al., 2002) and resistant regulatory genes (Mittler et al., 2004).

MDA content, final product of lipid peroxidation, reached higher values in *Fusarium* exposed plants of “Apache” than control plants on the 2nd day (Fig. 2B).

Our results are similar to research by El-Zahaby et al. (1995) who noted increased MDA content in resistant variety of barely after exposure to powdery mildew fungi. Furthermore, in “Apache”, MDA content showed decline after the 4th day of infection in regard to the 2nd day (Fig. 2B). Higher induction of POD activity (Fig. 1A), accompanied with the low MDA values (Fig. 2B) on the 4th day of treatment lead us to conclusion that POD in “Apache” plays an important role in maintaining the integrity of the membranes providing thus FHB-resistance. This is in accordance with the research by Zhang et al. (2013) where peroxidase increased remarkably in resistant wheat variety compared to susceptible one.

Conclusion

In “Apache”, *Fusarium* exposure caused the most notable changes in activity of the antioxidative enzymes. At an earlier time-point of infection (2nd day), the same genotype activates APX in response to *Fusarium*, while POD is a part of later defence strategies (4th day). Based on this, we can conclude that in “Apache”, APX and POD are important enzymes in providing the FHB-resistance. Unlike FHB-resistant “Apache”, treated plants of FHB-susceptible “Super Žitarka” sooner displayed the higher PPO activity (2nd day), which refers to the inclusion of non-enzymatic antioxidative as a reaction to *Fusarium* stress. In all tested genotypes, CAT did not show appreciable changes during *Fusarium* infection, which characterized it as a non-specific marker in screening the FHB disease.

Acknowledgments

This work has been supported in part by the Croatian Science Foundation within the project HRZZ-UIP-2014-9188.

References

- Aebi, H. (1984): Catalase in vitro, *Meth. Enzymol.* 105, 121-126. [https://doi.org/10.1016/s0076-6879\(84\)05016-3](https://doi.org/10.1016/s0076-6879(84)05016-3).
- Bai, G., Shaner, G. (2004): Management and resistance in wheat and barley to Fusarium head blight 1. *Annu. Rev. Phytopathol.* 42, 135-161. <https://doi.org/10.1146/annurev.phyto.42.040803.140340>.
- Boeckx, T., Winters, A. L., Webb, K. J., Kingston-Smith, A. H. (2015): Polyphenol oxidase in leaves; is there any significance to the chloroplastic localization? *J. Exp. Bot.* 1-9. <https://doi.org/10.1093/jxb/erv141>.
- Bradford, M. M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254. <https://doi.org/10.1006/abio.1976.9999>.
- Dat, J., Vandenaabeele, S., Vranova', E., Van Montagu, M., Inze', D., Van Breusegem, F. (2000): Dual action of the active oxygen species during plant stress responses. *Cell Mol. Life Sci.* 57, 779-795. <https://doi.org/10.1007/s000180050041>.
- El-Zahaby, H. M., Gullner, G., Kiraly, Z. (1995): Effects of powdery mildew infection of barley on the ascorbate-glutathione cycle and other antioxidants in different host-pathogen interactions. *Biochem. Cell Biol.* 85, 1225-1230. <https://doi.org/10.1094/phyto-85-1225>.
- Figueiró, A. A., Reese, N., Hernandez, J. L. G., Pacheco, M. T., Martinelli, J. A., Federizzi, L. C., Delatorre, C. A. (2015): Reactive oxygen species are not increased in resistant oat genotypes challenged by crown rust isolates. *J. Phytopathol.* 1-11. <https://doi.org/10.1111/jph.12377>.
- García-Limones, C., Hervás, A., Navas-Cortés, J. A., Jiménez-Díaz, R. M., Tena, M. (2002): Induction of an antioxidant enzyme system and other oxidative stress markers associated with compatible and incompatible interactions between chickpea (*Cicer arietinum* L.) and *Fusarium oxysporum* f. sp. ciceris. *Physiol. Mol. Plant Pathol.* 61, 325-337. <https://doi.org/10.1006/pmpp.2003.0445>.
- Gilbert, J., Tekauz, A. (2000): Review: recent developments in research on Fusarium head blight of wheat in Canada. *Can. J. Plant Pathol.* 22(1), 1-8. <https://doi.org/10.1080/07060660009501155>.
- Jain, A., Singh, A., Singh, S., Singh, H. B. (2013): Microbial consortium-induced changes in oxidative stress markers in pea plants challenged with *Sclerotinia sclerotiorum*. *J. Plant Growth Regul.* 32, 388-398. <https://doi.org/10.1007/s00344-012-9307-3>.
- Kuźniak, E., Skłodowska, M. (2005): Fungal pathogen-induced changes in the antioxidant systems of leaf peroxisomes from infected tomato plants. *Planta* 222, 192-200. <https://doi.org/10.1007/s00425-005-1514-8>.
- Kuźniak, E., Urbanek, H. (2000): The involvement of hydrogen peroxide in plant responses to stresses. *Acta Physiol. Plant.* 22(2), 195-203. <https://doi.org/10.1007/s11738-000-0076-4>.
- Li, W. M., Qian, C. M., Mo, Y. W., Hu, Y. L., Xie, J. H. (2011): Tolerance of banana for Fusarium wilt is associated with early H₂O₂ accumulation in the roots. *Afr. J. Biotechnol.* 10, 11378-11387. <https://doi.org/10.5897/ajmr11.298>.
- Madadkhah, E., Lotfi, M., Nabipour, A., Rahmanpour, S., Banihashemi, Z., Shooroei, M. (2012): Enzymatic activities in roots of melon genotypes infected with *Fusarium oxysporum* f. sp. melonis race 1. *Sci. Hortic.* 135, 171-176. <https://doi.org/10.1016/j.scienta.2011.11.020>.
- Mayer, A. M. (2006): Polyphenol oxidases in plants and fungi: going places? A review. *Phytochemistry* 67, 2318-2331. <https://doi.org/10.1016/j.phytochem.2006.08.006>.
- Mittler, R., Vanderauwera, S., Gollery, M., Van Breusegem, F. (2004): Reactive oxygen gene network of plant. *Trends Plant Sci.* 9, 490-498. <https://doi.org/10.1016/j.tplants.2004.08.009>.
- Mohammadi, M., Karr, A. (2002): α -1, 3-glucanase and chitinase activities in soybean root nodules. *J. Plant Physiol.* 159, 245-256. <https://doi.org/10.1078/0176-1617-00702>.
- Morkunas, I., Gemerek, J. (2007): The possible involvement of peroxidase in defence of yellow lupin embryo axes against *Fusarium oxysporum*. *Plant Physiol.* 164, 497-506. <https://doi.org/10.1016/j.jplph.2005.11.005>.
- Nakano, Y., Asada, K. (1981): Hydrogen peroxide is scavenged by ascorbate - specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 22, 867-880. <https://doi.org/10.1093/oxfordjournals.pcp.a076232>.
- Nanda, A. K., Andrio, E., Marino, D., Pauly, N., Dunand, C. (2010): Reactive oxygen species during plant-microorganism early interactions. *J. Integr. Plant Biol.* 52 (2), 195-204. <https://doi.org/10.1111/j.1744-7909.2010.00933.x>.
- Parveen, I., Threadgill, M. D., Moorby, J. M., Winters, A. (2010): Oxidative phenols in forage crops containing polyphenol oxidase enzymes. *J. Agric. Food Chem.* 58, 1371-1382. <https://doi.org/10.1021/jf9024294>.
- Racchi, M. L. (2013): Antioxidative defenses in plants with attention to *Prunus* and *Citrus* spp.. *Antioxidants* 2, 340-369. <https://doi.org/10.3390/antiox2040340>.
- Raymond, J., Rakariyatham, N., Azanza, J. (1993): Purification and some properties of polyphenol oxidase from sunflower seeds. *Phytochemistry* 34 (4), 927-931. [https://doi.org/10.1016/s0031-9422\(00\)90689-7](https://doi.org/10.1016/s0031-9422(00)90689-7).

- Shahbazi, H., Aminian, H., Sahebani, N., Halterman, D. A. (2010): Biochemical evaluation of resistance responses of potato to different isolates of *Alternaria solani*. *Phytopathology* 100, 454–459. <https://doi.org/10.1094/phyto-100-5-0454>.
- Siegel, B. Z., Galston, W. (1967): The isoperoxidases of *Pisum sativum*. *Plant Physiol.* 42, 221–226. <https://doi.org/10.1104/pp.42.2.221>.
- Sorahinobar, M., Niknam, V., Ebrahimzadeh, H., Soltanloo, H. (2015): Differential antioxidative responses of susceptible and resistant wheat cultivars against Fusarium head blight, *Intl. J. Farm. & Alli. Sci.* 4 (3), 239–243. <https://doi.org/10.1080/07060661.2017.1379442>.
- Španić, V., Viljevac Vuletić, M., Abičić, I., Marček, T. (2017): Early response of wheat antioxidant system with special reference to Fusarium head blight stress. *Plant Physiol. Biochem.* 115, 34–43. <https://doi.org/10.1016/j.plaphy.2017.03.010>.
- Thannickal, V. J., Fanburg, B. L. (2000): Reactive oxygen species in cell signalling. *Am. J. Physiol.* 279 (6), 1005–1028. https://doi.org/10.1007/0-306-48412-9_16.
- Thipyapong, P., Hunt, M. D., Steffens, J. C. (2004): Antisense downregulation of polyphenol oxidase results in enhanced disease susceptibility. *Planta* 220, 105–117. <https://doi.org/10.1007/s00425-004-1330-6>.
- Verma, S., Dubey, R. S. (2003): Leads toxicity induces lipid peroxidation and alters the activities of antioxidant enzymes in growing rice plants. *Plant Sci.* 164, 645–655. [https://doi.org/10.1016/s0168-9452\(03\)00022-0](https://doi.org/10.1016/s0168-9452(03)00022-0).
- Vranova, E., Inze, D., Van Breugsegem, F. (2002): Signal transduction during oxidative stress. *J. Exp. Bot.* 53, 1227–1236. <https://doi.org/10.1093/jexbot/53.372.1227>.
- Zadoks, J. C., Chang, T. T., Konzac, F. C. (1974): A decimal code for the growth stages of cereals. *Weed Res.* 14, 415–421. <https://doi.org/10.1111/j.1365-3180.1974.tb01084.x>.
- Zhang, P., Zhou, M. P., Zhang, X., Huo, Y., Ma, H. X. (2013): Change of defensive-related enzyme in wheat crown rot seedlings infected by *Fusarium graminearum*. *Cereal Res. Commun.* 41, 431–439. <https://doi.org/10.1556/crc.2013.0014>.